

# The *Ultimate* Formula Sheet for SAT Math

These formulas are provided in the reference information at the beginning of each SAT math section:

$$\text{Area of a Circle: } A = \pi r^2$$


$$\text{Circumference of a Circle: } C = 2\pi r$$

$$\text{Area of a Rectangle: } A = lw$$

$$\text{Area of a Triangle: } A = \frac{1}{2}bh$$

$$\text{Pythagorean Theorem: } a^2 + b^2 = c^2$$

Special Right Triangles:



$$\text{Volume of a Rectangular Prism (Box): } V = lwh$$

$$\text{Volume of a Cylinder: } V = \pi r^2 h$$

$$\text{Volume of a Sphere: } V = \frac{4}{3}\pi r^3$$

$$\text{Volume of a Cone: } V = \frac{1}{3}\pi r^2 h$$

$$\text{Volume of a Pyramid: } V = \frac{1}{3}lwh$$

**Fractions, Decimals, and Percentages: (for this section, r is the percent in decimal form)**

$$\text{Fraction} = \frac{\text{part}}{\text{whole}}$$

$$\text{percent} = \frac{\text{part}}{100}$$

Percent Increase or Decrease:

$$\frac{|\text{old} - \text{new}|}{\text{old}} \times 100\%$$

Increase by a percent: multiply by  $(1+r)$

Decrease by a percent: multiply by  $(1-r)$

$$\text{Simple Interest: } A = P(1+rt)$$

$$\text{Interest Compounded Annually: } A = P(1+r)^t$$

Interest Compounded n times per year:

$$A = P \left( 1 + \frac{r}{n} \right)^{nt}$$

**Rates, Ratios, and Proportions:**

General form of a conversion factor:

$$\left( \frac{\text{ending\_units}}{\text{starting\_units}} \right)$$

$$\text{Example: } 10\text{feet} \left( \frac{12\text{inches}}{1\text{foot}} \right) = 120\text{inches}$$

$$\begin{aligned} &\text{Concentration of A} \times \text{Volume of A} \\ &+ \text{Concentration of B} \times \text{Volume of B} \\ &= \text{Final concentration (Vol. of A + Vol. of B)} \end{aligned}$$

$$\text{Distance} = \text{Rate} \times \text{Time}$$

---

## Exponents, Roots, & Polynomials:

Multiplication Rule for Exponents:  $a^b \cdot a^c = a^{b+c}$

Division Rule for Exponents:  $\frac{a^b}{a^c} = a^{b-c}$

Power Rule for Exponents:  $(a^b)^c = a^{bc}$

Negative Exponents:  $a^{-b} = \frac{1}{a^b}$

Fractional Exponents:  $a^{\frac{b}{c}} = \sqrt[c]{a^b}$  or  $(\sqrt[c]{a})^b$

$i^2 = -1$ ;  $i^3 = -i$ ;  $i^4 = 1$

$i^{4n} = 1$ ;  $i^{4n+1} = i$ ;  $i^{4n+2} = -1$ ;  $i^{4n+3} = -i$

---

## Parabolas:

Standard Form:  $f(x) = ax^2 + bx + c$ ;

vertex =  $\left(-\frac{b}{2a}, f\left(-\frac{b}{2a}\right)\right)$ ;

y-intercept = c;

x-intercepts =  $\frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

Sum of solutions =  $-\frac{b}{a}$

Discriminant =  $b^2 - 4ac$ ; Pos=2 real roots Zero=1 real root; Neg=2 imaginary roots


Factored Form:  $f(x) = a(x - m)(x - n)$ ;

x-intercepts are m and n;

x-coordinate of vertex =  $\frac{m+n}{2}$

Vertex Form:  $f(x) = a(x - h)^2 + k$ ;

vertex =  $(h, k)$



Difference of Squares:  $a^2 - b^2 = (a+b)(a-b)$

Perfect Square Trinomial:  $a^2 + 2ab + b^2 = (a+b)^2$  and  $a^2 - 2ab + b^2 = (a-b)^2$

Completing the Square:  $x^2 + bx + \left(\frac{b}{2}\right)^2 = \left(x + \frac{b}{2}\right)^2$

---


---

### Graphing Lines:

$$\text{Slope Formula: } m = \frac{y_2 - y_1}{x_2 - x_1}$$

Slope of horizontal line = 0

Slope of vertical line = undefined



$$\text{Standard Form: } Ax + By = C$$

$$\text{Slope-Intercept Form: } y = mx + b$$

$$\text{Point-Slope Form: } y - y_1 = m(x - x_1)$$

$$\text{Distance Formula: } d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

$$\text{Midpoint Formula: } M = \left( \frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2} \right)$$

Parallel lines: equal slopes

⊥ Lines: slopes are opposite reciprocals

---

### Data and Probability:

$$\text{average} = \frac{\text{sum of items}}{\text{number of items}}$$

median = middle number

range = maximum - minimum

$$\text{probability} = \frac{\text{desired outcomes}}{\text{possible outcomes}}$$

---

### Angles:

Vertical ∠'s are ≅

∠'s that form a linear pair are supplementary (add up to 180°)

∠'s that form a circle add up to 360°

When ∥ lines are cut by a transversal, all acute ∠'s are ≅ and all obtuse ∠'s are ≅

### Triangles:


The three ∠'s of a Δ add up to 180°

An exterior ∠ is equal to the sum of the two remote interior ∠'s


Pythagorean Triples: 3-4-5 and 5-12-13

### Circles:

A radius and tangent make a right ∠



A central ∠ is double the inscribed ∠



$$\frac{x}{360} = \frac{\text{arc}}{\text{circumference}} \quad \text{and} \quad \frac{x}{360} = \frac{\text{sector}}{\text{area of circle}} \quad \text{where } x = \text{central angle}$$

Formula for a Circle:  $(x - h)^2 + (y - k)^2 = r^2$ , where (h,k) is the center and r is the radius

**Polygons: (for this section, n is the number of sides)**

Area of a trapezoid:  $\frac{1}{2}(b_1 + b_2)h$

One interior angle of a regular polygon:  
 $\frac{180(n-2)}{n}$

Sum of the interior angles:  $180(n-2)$

Sum of the exterior angles:  $360^\circ$

**Properties of Parallelograms:**


1. Opp sides are  $\parallel$  and  $\cong$
2. Opp  $\angle$ 's are  $\cong$
3. Consec  $\angle$ 's are supplementary
4. Each diagonal forms a pair of  $\cong \Delta$ 's
5. Diagonals bisect each other  
→ If they are  $\cong$  it is a rectangle  
→ If they are  $\perp$  it is a rhombus
6. *Area = base  $\times$  height*

**Trigonometry:**

$\sin = \frac{opp}{hyp}$      $\cos = \frac{adj}{hyp}$      $\tan = \frac{opp}{adj}$      $360^\circ = 2\pi$  radians

$\sin(x) = \cos(90 - x)$  The sine of an  $\angle$  is equal to the cosine of its complement.

**Parent Graphs & Transformations:**



**Transformation**

$f(x) + k$

$f(x) - k$

$f(x + h)$

$f(x - h)$

$-f(x)$

$cf(x)$

$\frac{1}{c}f(x)$

**Visual effect**

Shift up by k units

Shift down by k units

Shift left by h units

Shift right by h units

Reflect over the x axis (flip upside down)

Stretch vertically by a factor of c (becomes skinnier)

Shrink vertically by a factor of c (becomes fatter)